

Montana Bureau of Mines and Geology
Open File No. 483

Preliminary Geologic Map of the Philipsburg
30' x 60' Quadrangle, Western Montana

Jeffrey D. Lonn, Catherine McDonald, Reed S. Lewis, Thomas J. Kalaka
J. Michael O'Neill, Richard B. Berg, and Phyllis Hargrave

2003

To view a full scale version of this map, [click here](#).


For the text files with the map information, [click here](#).

[Digital data link](#)

Note — This map was originally published at a scale of 1:100,000 but the page sizes have been modified to fit average printer capabilities (8½ x 14; legal size paper). There is an eighth inch overlap on these pages. A full sized colored print of this map can be ordered from the MBMG Publication Sales Office, 1300 West Park Street, Butte, MT, 59701-8997.

Phone: 406-496-4167

Email: pubsales@mbmg.mtech.edu



Montana Bureau of Mines and Geology
Open File 483 Plate 1

**Geologic Map of the
Philipsburg 30' x 60' Quadrangle,
Western Montana**

Jeffrey D. Lonn¹, Catherine McDonald¹, Reed S. Lewis²,
Thomas J. Kalakay³, J. Michael O'Neill⁴,
Richard B. Berg¹, and Phyllis Hargrave¹

2003 (Revised 2010)

¹ Montana Bureau of Mines and Geology; ² Idaho Geological Survey;
³ Rocky Mountain College; ⁴ U.S. Geological Survey



Quad index

MISSOULA WEST	MISSOULA EAST	ELLISTON
HAMILTON	PHILIPSBURG	BUTTE NORTH
NEZ PERCE PASS	WISDOM	BUTTE SOUTH


Partial support has been provided by the STATEMAP component of the National Cooperative Geologic Mapping Program of the U.S. Geological Survey under Contract Number 02 HQAG0038.



GIS production: Ken Sandau and Paul Thale, MBMG. Map layout: Susan Smith, MBMG.

Revised	Date
Map and text by Jeffrey D. Lonn	6/09
Map and text by Jeffrey D. Lonn	7/10



Maps may be obtained from
Publications Office
Montana Bureau of Mines and Geology
1300 West Park Street, Butte, Montana 59701-8997
Phone: (406) 496-4167 Fax: (406) 496-4451
<http://www.mbg.mtech.edu>





SOURCES OF PREVIOUS GEOLOGIC MAPPING

- | | | | |
|--|-----------------------------|--|------------------------------|
| | 1. Allen, 1961 | | 16. Loen and others, 1989 |
| | 2. Bakken, 1984 | | 17. Lofholm, 1985 |
| | 3. Buckley, 1990 | | 18. Mahorney, 1956 |
| | 4. Csejtey, 1962 | | 19. McGill, 1961 |
| | 5. Desmarais, 1983 | | 20. Mutch, 1961 |
| | 6. Earll, 1972 | | 21. Noel, 1956 |
| | 7. Elliott and others, 1984 | | 22. Pederson, 1976 |
| | 8. Emmons and Calkins, 1913 | | 23. Poulter, 1956 |
| | 9. Heise, 1983 | | 24. Presley, 1971 |
| | 10. Hughes, 1970 | | 25. Prinz, 1967 |
| | 11. Langton, 1935 | | 26. Stuart, 1966 |
| | 12. LaTour, 1974 | | 27. Wallace, 1987 |
| | 13. Lelek, 1979 | | 28. Wallace and others, 1986 |
| | 14. Lewis, 1998a | | 29. Wallace and others, 1989 |
| | 15. Lidke and Wallace, 1992 | | 30. Wallace and others, 1992 |
- These sources cover the whole 100,000 scale quadrangle




Figure 1. Major tectonic features, structural domains, and mountain ranges of the Philipsburg 30' x 60' quadrangle.

MAP UNIT DESCRIPTIONS

<div style="background-color: #ffff00; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qmd</p>	<p>MINE WASTE (HOLOCENE) Piles of poorly sorted cobbles, boulders, and sand resulting from placer mining operations. Thickness 1.5-7.5 m (5-25 ft).</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qc</p>	<p>COLLUVIUM (HOLOCENE) Thin, unconsolidated slope wash, talus, and rock fall deposits. Thickness 1.5-7.5 m (5-25 ft).</p>
<div style="background-color: #ffff00; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qta</p>	<p>TALUS DEPOSITS (HOLOCENE AND PLEISTOCENE) Accumulations of angular boulders below cliffs. Thickness 1.5-15.0 m (5-25 ft).</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qal</p>	<p>ALLUVIUM OF MODERN CHANNELS AND FLOODPLAINS (HOLOCENE) Mostly well-rounded, well-sorted boulders, cobbles, gravel, sand, and silt deposited in modern stream channels and floodplains. Includes both fine-grained overbank deposits and coarse-grained channel deposits. In some areas, older alluvium (Qao) is not divided from Qal.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qao</p>	<p>OLDER ALLUVIUM (EARLY HOLOCENE AND LATE PLEISTOCENE?) Mostly well-rounded, well-sorted boulders, cobbles, sand, and silt deposited by streamflow processes. Surfaces of these deposits now stand 1.5-12.1 m (5-40 ft) above the modern floodplain. Includes terrace deposits along streams and glacial outwash deposited by braided streams.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qaf</p>	<p>ALLUVIAL FAN DEPOSITS (HOLOCENE AND PLEISTOCENE) Poorly to well-sorted, rounded to sub-angular boulders, cobbles, sand, silt, and clay. Surfaces of these deposits have a distinct fan shape. Deposited by both stream flow and debris flow processes in alluvial fan environments. In some areas, older alluvial fan deposits (Qafo) are not divided from Qaf.</p>
<div style="background-color: #ffff00; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qafo</p>	<p>OLDER ALLUVIAL FAN DEPOSITS (EARLY HOLOCENE AND LATE PLEISTOCENE?) Fans whose surfaces are now perched 1.5-12.1 m (5-40 ft) above the modern landforms.</p>
<div style="background-color: #ffff00; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qrg</p>	<p>ROCK GLACIER DEPOSITS (HOLOCENE) Lobate accumulations of angular boulders emplaced by flow of an ice core. Active and inactive rock glaciers are not divided.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qls</p>	<p>LANDSLIDE DEPOSITS (HOLOCENE AND PLEISTOCENE) Unsorted and unstratified mixtures of mud and boulders transported by mass movement down steep slopes. Characterized by irregular topography.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qdf</p>	<p>DEBRIS FLOW DEPOSITS (PLEISTOCENE) Poorly sorted, sub-angular bouldery deposits of huge boulders, cobbles, sand, silt and clay deposited by catastrophic debris flows.</p>
<div style="background-color: #ffff00; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qgtk</p>	<p>GLACIAL TILL AND KAME DEPOSITS, UNDIVIDED (PLEISTOCENE)</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qgt</p>	<p>GLACIAL TILL (PLEISTOCENE) Unsorted, mostly unstratified, clay, silt, sand, and gravel with subrounded boulders as much as 3 m (10 ft) in diameter. Till is often characterized by large, subrounded, exotic boulders that have been transported some distance, and by hummocky topography. Poor drainage, with swampy areas and numerous springs, and subangular clasts distinguish it in the field from kame deposits (Qgk).</p>
<div style="background-color: #ffff00; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Qgk</p>	<p>GLACIAL KAME DEPOSITS (PLEISTOCENE) Moderately to well-sorted, sub-rounded to well-rounded, well-stratified sand, pebbles, and boulders deposited by streams flowing within, on, and marginal to glaciers. Topographic surfaces</p>

<div style="background-color: #ffff00; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Kqd</p>	<p>QUARTZ DIORITE (LATE CRETACEOUS) Hornblende-biotite quartz diorite, diorite, and granodiorite.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Kgd</p>	<p>GRANODIORITIC ROCKS (LATE CRETACEOUS) Biotite-hornblende granodiorite, biotite granodiorite, and tonalite.</p>
<div style="background-color: #ffff00; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Kgdp</p>	<p>PORPHYRITIC GRANODIORITE (LATE CRETACEOUS) Porphyritic, muscovite-bearing granodiorite of the Mount Powell Batholith.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Kgdf</p>	<p>FOLIATED GRANODIORITE (LATE CRETACEOUS) Moderately to strongly foliated biotite-hornblende granodiorite.</p>
<div style="background-color: #ffff00; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Kqdf</p>	<p>FOLIATED QUARTZ DIORITE (LATE CRETACEOUS) Weakly to strongly foliated quartz diorite of the eastern Anaconda Range.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Kcg</p>	<p>COLORADO GROUP (EARLY TO LATE CRETACEOUS) The upper Colorado Group consists of approximately 300 m (1,000 ft) of tan to brown lithic sandstone, gray to gray-green siltstone, minor shale, and local conglomerate. Sandstones often contain dark chert grains or volcanic fragments. The lower Colorado Group includes dark gray to black fissile shale underlain by tan to gray siltstone, fine-grained sandstone, and dark gray to black limestone. This lower part is more calcareous, and limestone is more abundant near the gradational contact with the underlying Kootenai Formation. The thickness of the Colorado Group is about 498 m (1,635 ft) (McGill, 1961).</p>
<div style="background-color: #ffff00; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Kcgp</p>	<p>PHYLLITE OF THE METAMORPHOSED COLORADO GROUP (EARLY TO LATE CRETACEOUS) Phyllite, quartzite, and minor marble that are the metamorphic equivalent of the Colorado Group.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>KJs</p>	<p>SEDIMENTARY ROCKS OF THE ELLIS AND KOOTENAI FORMATIONS, UNDIVIDED (JURASSIC AND CRETACEOUS) The upper part of the non-marine Kootenai Formation includes as much as 40 m (130 ft) of gray, gastropod-bearing limestone at the top underlain by gray to reddish calcareous shale, siltstone, and sandstone. The middle part consists of about 275 m (900 ft) of maroon, green, and gray mudstone and siltstone with minor calcareous sandstone. The lower Kootenai, approximately 91 m (300 ft) thick, includes an interval of finely crystalline gray limestone with varicolored siltstone and mudstone mostly underlain by gray to red-brown feldspathic sandstone and siltstone with widespread gray-chert conglomerate. The upper Ellis Formation is calcareous gray sandstone, siltstone, and shale with locally thin, calcareous conglomeratic beds containing abundant pebbles and sand-sized grains of black chert. The lower Ellis includes non-calcareous black fissile shales, dark gray to black calcareous shales and siltstones, and thin limestone fossil-hash beds. The Ellis Formation is approximately 85 m (280 ft) thick.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>KJsp</p>	<p>PHYLLITE AND QUARTZITE OF THE METAMORPHOSED KOOTENAI AND ELLIS FORMATIONS (JURASSIC AND CRETACEOUS) Phyllite, quartzite, marble, and schist that are the metamorphic equivalents of the Kootenai and Ellis Formations.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>PIPs</p>	<p>SEDIMENTARY ROCKS OF THE SHEDHORN, PHOSPHORIA, PARK CITY AND QUADRANT FORMATIONS, UNDIVIDED (PERMIAN AND PENNSYLVANIAN) The Shedhorn Formation is tan to white orthoquartzite, locally cross-bedded and pitted, approximately 9 m (30 ft) thick. The Phosphoria Formation is bedded, dark gray to red chert, chert conglomerate, minor calcareous siltstone, gray to black phosphatic shale, phosphorite, and argillaceous carbonate that is commonly oolitic. Approximately 20 m (65 ft) thick. The Park City Formation is predominantly calcareous sandstone, siltstone, and nodular cherty limestone</p>

<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Ygr</p>	<p>GARNET RANGE FORMATION (MESOPROTEROZOIC) Greenish-gray, micaceous, tabular and lenticular, hummocky quartzite with argillite interbeds. In some areas, purple and white quartzite with argillite interbeds. Mesoproterozoic Pilcher Formation overlies the Garnet Range Formation. Much of the map area, but it has a thickness of about 305 m (1,000 ft) near Georgetown Lake.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Ym</p>	<p>MCNAMARA FORMATION (MESOPROTEROZOIC) Beds of flat-laminated and trough crossbedded, fine- to medium-grained argillite and thin argillite beds. Cherty rip-up clasts are common and diagnostic in its type locality near Missoula. From 0 to 457 m (1,500 ft) thick.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Ybo</p>	<p>BONNER FORMATION (MESOPROTEROZOIC) Pink, medium-grained, feldspathic, crossbedded quartzite. In some areas, pebbles are abundant. Lewis (1998b) found 15-25% potassium feldspar and plagioclase in the Bonner Formation. Approximately 518 m (1,700 ft) thick.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Yboq</p>	<p>QUARTZITE OF THE METAMORPHOSED BONNER FORMATION (MESOPROTEROZOIC) Highly recrystallized quartzite with minor phyllite and schist. Metamorphic equivalent of the Bonner Formation.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Yms</p>	<p>MOUNT SHIELDS FORMATION (MESOPROTEROZOIC) Usually divided into 3 informal members that are not distinguished in the field. Total thickness is approximately 1,219 m (4,000 ft).</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Yms3</p>	<p>MOUNT SHIELDS FORMATION, MEMBER 3, INFORMAL (MESOPROTEROZOIC) Mostly red siltite to argillite couples and couplets with abundant pebbles and clasts.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Yms2</p>	<p>MOUNT SHIELDS FORMATION, MEMBER 2, INFORMAL (MESOPROTEROZOIC) Pink to gray, flat-laminated, fine- to medium grained quartzite with argillite blebs. Contains some crossbeds. In the Anaconda Range of the western Anaconda Range, and in the Skalkaho region, the Mount Shields contains a variety of lithologies making it difficult to distinguish from the Bonner Formation. Lewis (1998b) found subequal amounts of plagioclase and potassium feldspar content of 25-35% in the Mount Shields.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Ymsq</p>	<p>QUARTZITE OF THE METAMORPHOSED MOUNT SHIELDS FORMATION (MESOPROTEROZOIC) Metamorphic equivalent of the Mount Shields Member 3.</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Yss</p>	<p>SHEPARD AND SNOWSLIP FORMATIONS, UNDIVIDED (MESOPROTEROZOIC) Total thickness as much as 1,067 m (3,500 ft).</p>
<div style="background-color: #d3d3d3; border: 1px solid black; width: 20px; height: 10px; margin-bottom: 5px;"></div> <p>Ysh</p>	<p>PHYLLITE, QUARTZITE, AND CALC-SILICATE ROCKS OF THE METAMORPHOSED SNOWSLIP FORMATIONS (MIDDLE PROTEROZOIC) Metamorphic equivalent of these formations.</p>

REFERENCES

- Allen, J.C., 1961, Structure and petrology of the Royal Stock and the Mount Powell Batholith, Flint Creek Range, western Montana: Princeton, NJ, Princeton University, Ph.D. dissertation, 112 p., map scale 1:62,500.
- Bakken, J.F., 1984, The structural geology and tectonic history of the northern Flint Creek Range, western Montana: Bozeman, Montana State University, M.S. thesis, 125 p.
- Berg, R.B., and Dahy, J.P., 2002, Montana sapphires and speculation on their origin, in Scott, P.W., and Bristow, C.M., eds., Industrial Minerals and Extractive Industry Geology: Geological Society of London, p. 199-204.
- Bickford, M.E., Chase, R.B., Nelson, B.K., Schuster, R.D., and Arruda, E.C., 1981, U-Pb studies of zircon cores and overgrowths, and monazite-implications for age and petrogenesis of the northeastern Idaho Batholith: *Journal of Geology*, v. 89, p. 433-547.
- Buckley, S.N., 1990, Ductile extension in a Late Cretaceous fold and thrust belt, Granite County, west-central Montana: Missoula, University of Montana, M.S. thesis, 42 p., map scale 1:6,000.
- Burchfiel, B.C., and Royden, L.H., 1985, North-south extension within the convergent Himalayan region: *Geology*, v. 13, p. 679-682.
- Chase, R.B., Bickford, M.E., and Arruda, E.C., 1983, Tectonic implications of Tertiary intrusion and shearing within the Bitterroot dome, northeastern Idaho Batholith: *Journal of Geology*, v. 91, p. 462-470.
- Clark, S.L., 1979, Structural and petrological comparison of the southern Sapphire Range, Montana, with the northeast border zone of the Idaho Batholith: Kalamazoo, Western Michigan University, M.S. thesis, 88 p., map scale 1:48,000.
- Csejtey, Bela, 1962, Geology of the southeast flank of the Flint Creek Range, western Montana: Princeton, NJ, Princeton University, Ph.D. dissertation, 175 p., map scale 1:62,500.
- Culshaw, N.G., Beaumont, C., and Jamieson, R.A., 2006, The orogenic superstructure-infrastructure concept: Revisited, quantified, and revived: *Geology*, v. 34, p. 733-736.
- Dalmayrac, Bernard, and Molnar, Peter, 1981, Parallel thrust and normal faulting in Peru and constraints on the state of stress: *Earth and Planetary Science Letters*, v. 55, p. 473-481.
- Desmarais, N.R., 1983, Geology and geochronology of the Chief Joseph plutonic-metamorphic complex, Idaho-Montana: Seattle, University of Washington, Ph.D. dissertation, 150 p., map scale 1:48,000.
- Doughty, P.T., and Sheriff, S.D., 1992, Paleomagnetic evidence for an echelon crustal extension and crustal rotations in western Montana and Idaho: *Tectonics*, v. 11, p. 663-671.
- Earll, F.N., 1972, Mines and mineral deposits of the southern Flint Creek Range, western Montana: Bozeman, Montana State University, M.S. thesis, 125 p., map scale 1:62,500.
- Hyndman, D.W., Silverman, A.J., Ehinger, R., and Benoit, W., 1982, The Philipsburg Batholith, western Montana: mineralogy, petrology, internal variation, and evolution: *Montana Bureau of Mines and Geology Memoir M 49*, 37 p.
- Hyndman, D.W., Alt, David, and Sears, J.W., 1988, Post-Archean metamorphic and tectonic evolution of western Montana and northern Idaho, in Ernst, W.G., ed., *Metamorphism and Crustal Evolution in the Western Conterminous U.S.* (Rubey Volume VII): Englewood Cliffs, NJ, Prentice-Hall, p. 332-361.
- Kalakay, T.J., and Lonng, J.D., 2002, Geometric and kinematic relationships between high-temperature and low-temperature faulting in the Anaconda detachment zone, southwest Montana: *Geological Society of America Abstracts with Programs*, v. 34, no. 6, p. 370.
- Kalakay, T.J., Foster, D.A., and Thomas, R.C., 2003, Geometry and timing of deformation in the Anaconda extensional terrane, west-central Montana: *Northwest Geology*, v. 32, p. 124-133.
- Langton, C.M., 1935, Geology of the northeastern part of the Idaho Batholith and adjacent region in Montana: *Journal of Geology*, v. 43, p. 35-60, map scale 1:422,000.
- LaTour, T.E., 1974, An examination of metamorphism and scapolite in the Skalkaho region, southern Sapphire Range: Missoula, University of Montana, M.S. thesis, 95 p.
- Lelek, J.J., 1979, The Skalkaho pyroxenite-syenite complex east of Hamilton, Montana, and the role of magma immiscibility in its formation: Missoula, University of Montana, M.S. thesis, 130 p.
- Lewis, R.S., 1998a, Geologic map of the Butte 10 x 20 quadrangle: Montana Bureau of Mines and Geology Open File Report MBMG 363, scale 1:250,000.
- Lewis, R.S., 1998b, Stratigraphy and structure of the lower Missoula Group in the Butte 1° x 2° and Missoula West 30' x 60' quadrangles, Montana: *Northwest Geology*, v. 28, p. 1-14.
- Lidke, D.J., and Wallace, C.A., 1992, Rocks and structure of the north-central part of the Anaconda Range, Deer Lodge and Granite Counties, Montana: *U.S. Geological Survey Bulletin* 1993, 31 p., map scale 1:24,000.
- Loen, J.S., Blaskowski, M.J., and Elliott, J.E., 1989, Geology and mineral deposits of the Miner's Gulch area, Granite County, Montana: *U.S. Geological Survey Bulletin* 1791, 51 p., map scale 1:10,000.
- Lofholm, S.T., 1985, Geology of the Old Dominion Mine prospect, Granite County, Montana: Rapid City, South Dakota School of Mines, M.S. thesis, 52 p., map scale 1:6,000.
- Lonng, J.D., and Berg, R.B., 1999, Preliminary geologic map of the Hamilton Range, western Montana: Missoula, Montana Bureau of Mines and Geology Open File Report MF-1924, scale 1:250,000.
- Reid, R.R., Wavra, C.S., and Bond, W.D., 1995, Constriction fracture flow: A mechanism for fault and vein formation in the Coeur d'Alene district, Idaho: *Economic Geology*, v. 90, p. 81-87.
- Ruppel, E.T., O'Neill, J.M., and Lopez, D.A., 1993, Geologic map of the Dillon 1° x 2° quadrangle, Idaho and Montana: *U.S. Geological Survey Miscellaneous Investigations Series Map I-1803-H*, scale 1:250,000.
- Stuart, C.J., 1966, Metamorphism in the central Flint Creek Range, Montana: Missoula, University of Montana, M.S. thesis, 103 p., map scale 1:6,000.
- Tysdal, R.G., Hanna, W.F., and Capstick, D.O., 1988, Mineral resources of the Quigg West Wilderness Study Area, Granite County, Montana: *U.S. Geological Survey Bulletin* 1724-D, 25 p., map scale 1:24,000.
- Wallace, C.A., 1987, Generalized geologic map of the Butte 10 x 20 quadrangle, Montana: *U.S. Geological Survey Miscellaneous Field Studies Map MF-1924*, scale 1:250,000.
- Wallace, C.A., Schmidt, R.G., Lidke, D.J., Waters, M.R., Elliott, J.E., French, A.B., Whipple, J.W., Zarske, S.E., Blaskowski, M.J., Heise, B.A., Yeoman, R.A., O'Neill, J.M., Lopez, D.A., Robinson, G.D., and Klepper, M.R., 1986, Preliminary geologic map of the Butte 1° x 2° quadrangle, Montana: *U.S. Geological Survey Open File Report* 86-292, scale 1:250,000.
- Wallace, C.A., Lidke, D.J., Waters, M.R., and Obradovich, J.D., 1989, Rocks and structure of the southern Sapphire Mountains, Granite and Ravalli Counties, western Montana: *U.S. Geological Survey Bulletin* 1824, 29 p., map scale 1:50,000.
- Wallace, C.A., Lidke, D.J., Elliott, J.E., Desmarais, N.R., Obradovich, J.D., Lopez, D.A., Zarske, S.E., Heise, B.A., Blaskowski, M.J., and Loen, J.F., 1992, Geologic map of the Anaconda-Pintlar Wilderness and contiguous roadless areas, Granite, Deerlodge, Beaverhead, and Ravalli Counties, western Montana: *U.S. Geological Survey Miscellaneous Field Studies Map MF-1633C*, 36 p., scale 1:50,000.
- Winegar, R.C., 1971, The petrology of the Lost Creek stock and its relation to the Mount Powell Batholith, Montana: Missoula, University of Montana, M.S. thesis, 60 p.
- Winston, Don, 1986a, Sedimentology of the Ravalli Group, middle Belt carbonate and Missoula Group, middle Proterozoic Belt Supergroup, Montana, Idaho, and Washington, in Roberts, S.M., *Belt Supergroup: A guide to Proterozoic rocks of western Montana and adjacent areas*: Montana Bureau of Mines and Geology Special Publication SP 94, p. 85-124.
- Winston, Don, 1986b, Middle Proterozoic tectonics of the Belt Basin, western Montana and northern Idaho, in Roberts, S.M., ed., *Belt Supergroup: A guide to Proterozoic rocks of western Montana and adjacent areas*: Montana Bureau of Mines and Geology Special Publication SP 94, p. 245-257.

QTaf	ALLUVIAL FAN DEPOSITS (EARLY PLEISTOCENE AND LATE TERTIARY) Poorly to well-sorted, rounded to sub-angular boulders, cobbles, sand, silt, and clay. Surfaces of these deposits have a distinct fan shape and now stand more than 15 m (50 ft) above modern deposits.
Ts	SEDIMENTARY ROCKS, UNDIVIDED (TERTIARY) Include both coarse- and fine-grained rocks.
Taf	ALLUVIAL FAN DEPOSITS (TERTIARY) Poorly to well-sorted, rounded to sub-angular boulders, cobbles, sand, silt, and clay. Surfaces of these deposits now stand more than 15 m (50 ft) above modern deposits. Unlike younger alluvial fan deposits (QTaf), these are often unrelated to modern drainage patterns.
Tsc	SIXMILE CREEK FORMATION (PLIOCENE AND MIOCENE?) Mostly conglomerate with some sandstone and siltstone. Commonly caps the remnant Tertiary surfaces.
Tcl	CLAY AND SILT (MIOCENE AND OLIGOCENE?) White to light-gray clay and silt deposited in fluvial and lacustrine environments, and probably correlative with the Renova Formation.
Tac	ANACONDA BEDS, INFORMAL (EOCENE?) Unstratified deposits of angular, poorly sorted boulders and cobbles of unmetamorphosed sedimentary rocks deposited in debris flow and mass wasting environments, interbedded with and grading to moderately sorted, sub-angular to rounded cobbles, pebbles, and sand deposited in fluvial environments. Emmons and Calkins (1913) first described these rocks in the Barker Creek area southeast of Silver Lake and called them "earlier Tertiary gravels". Csejtey (1962) described similar rocks near Anaconda and named them the Anaconda beds. They have been interpreted as deposits formed during the unroofing of the Anaconda core complex in basins developed along the detachment zone (O'Neill, 2005).
Tcg	CONGLOMERATE (EOCENE?) Clast-supported, sub-angular to rounded, moderately sorted boulders, cobbles, and pebbles of Belt Supergroup rocks. Contains rare volcanic clasts. Found in the area of the Rock Creek volcanic field.
Tv	VOLCANIC ROCKS, UNDIVIDED (TERTIARY) Volcanic and volcanoclastic rocks. Include minor hypabyssal intrusive bodies.
Ttb	TUFF BRECCIA (EOCENE?) Poorly stratified, poorly sorted, mostly clast-supported, angular boulder- to cobble-sized lithic fragments in an ash matrix. Clasts are predominantly local Belt Supergroup sedimentary rocks, but locally include abundant volcanic rocks. Matrix material is usually sparse, although in some areas ash-flow tuff predominates with few lithic fragments. Found in the area of the Rock Creek volcanic field.
Tr	RHYOLITE (EOCENE?) Rhyolite flows and tuff that contain abundant biotite phenocrysts and sparse potassium feldspar and quartz phenocrysts. Mapped in the Rock Creek volcanic field.
Tri	INTRUSIVE RHYOLITE (EOCENE?) Dikes containing euhedral potassium feldspar phenocrysts as much as 15 mm (0.5 in) long. Also contains sparse plagioclase, biotite, and quartz phenocrysts.
Tlc	LOWLAND CREEK VOLCANICS (EOCENE) Rhyolite and dacite flows, tuffs, and volcanoclastic rocks.
Trt	RHYOLITE TUFF (EOCENE) Mapped within the Lowland Creek volcanic field.
Tbmg	BIOTITE-MUSCOVITE GRANITE (EOCENE AND PALEOCENE) Equigranular and porphyritic biotite-muscovite granite. Mylonitic foliation is present within and adjacent to the Anaconda detachment fault.
Tgd	GRANODIORITIC ROCKS (EOCENE AND PALEOCENE) Quartz monzodiorite and granodiorite.

PPsq	QUARTZITE OF THE METAMORPHOSED SHEDHORN, PHOSPHORIA, PARK CITY AND QUADRANT FORMATIONS (PERMIAN AND PENNSYLVANIAN) Mostly quartzite, but also includes minor phyllite and marble. Metamorphic equivalent of the Shedhorn, Phosphoria, Park City, and Quadrant Formations.
PMs	SEDIMENTARY ROCKS OF THE SNOWCREST RANGE GROUP, AMSDEN FORMATION, AND MADISON GROUP, UNDIVIDED (PENNSYLVANIAN AND MISSISSIPPIAN) The Snowcrest Range Group (Lidke and Wallace, 1992) and equivalent Amsden Formation (Emmons and Calkins, 1913; Poultter, 1956; McGill, 1961) consist mainly of maroon, thin-bedded dolomite overlain by maroon dolomitic shale with minor light gray limestone and dolomite. Recessive weathering; generally mapped by presence of red soil zone. Approximately 91 m (300 ft) thick. The Madison Group consists of massive-weathering, thick-bedded, white to bluish-gray, fossiliferous, cherty limestone underlain by dark gray, flaggy limestone with interbedded, black calcareous shale. Approximate thickness as much as 1,022 m (2,300 ft) (McGill, 1961).
IPMsm	MARBLE OF THE METAMORPHOSED SNOWCREST RANGE GROUP, AMSDEN FORMATION, AND MADISON GROUP (PENNSYLVANIAN AND MISSISSIPPIAN) Mostly marble, but also includes some minor phyllite. Metamorphic equivalent of the Snowcrest Range Group, Amsden Formation, and Madison Group. This unit is often tectonically thinned. See geologic discussion for more detail.
MDsm	MARBLE OF METAMORPHOSED SEDIMENTARY ROCKS, UNDIVIDED (MISSISSIPPIAN AND DEVONIAN) Metamorphic equivalents of Madison Group, Three Forks Formation, Jefferson Dolomite, and Maywood Formation.
MDsm	MARBLE OF METAMORPHOSED SEDIMENTARY ROCKS, UNDIVIDED (MISSISSIPPIAN AND DEVONIAN) Metamorphic equivalents of Madison Group, Three Forks Formation, Jefferson Dolomite, and Maywood Formation.
Ds	SEDIMENTARY ROCKS OF JEFFERSON AND MAYWOOD FORMATIONS, UNDIVIDED (UPPER DEVONIAN) The Jefferson Formation, approximately 260 m (850 ft) thick, consists of thick-bedded, dark gray to black dolomite with minor interbedded light gray limestone. Limestone beds are more common in upper part and are often brecciated. Alternating light and dark beds and a petroliferous odor are characteristic. The Maywood Formation consists of thin bedded, gray, reddish gray, and yellow dolomitic shale and siltstone, silty dolomite, and sparse gray limestone. Upper part contains minor quartzite and dark dolomite beds similar to overlying Jefferson Formation. Basal part contains beds of dolomitic and calcareous sandstone and siltstone. Thickness 84-106m (275-350 ft).
Dsm	MARBLE OF THE METAMORPHOSED JEFFERSON AND MAYWOOD FORMATIONS (UPPER DEVONIAN) Includes marble that is the metamorphic equivalent of the Jefferson Formation and quartzite and phyllite equivalent to the Maywood Formation. This unit is often tectonically thinned. See geologic discussion for more detail.
Cs	SEDIMENTARY ROCKS OF THE RED LION, HASMARK, SILVER HILL AND FLATHEAD FORMATIONS, UNDIVIDED (CAMBRIAN) The Red Lion Formation is predominately light gray to blue gray limestone with yellow- to red-weathering, wavy, dolomitic and siliceous laminae. Laminae are less common near top of formation. Basal part contains black calcareous shale with minor sandstone and limestone. The thickness is approximately 100 m (330 ft). The Hasmark Formation, approximately 320 m (1,050 ft) thick, is a uniform, light to blue gray dolomite that is separated into an upper and lower part by a recessive shale interval. The lower dolomite is generally darker in color than the upper part and commonly contains oolitic structures and mottled weathered surfaces. The shale interval is as much as 46 m (150 ft) thick and varies from dark brown to reddish purple calcareous shale to reddish limestone and shale. The Silver Hill Formation, approximately 185 m (600 ft) thick, includes calcareous brown, white, and green shale interbedded with laminated limestone underlain by moderately thick-bedded, laminated, light-gray limestone. Laminae are generally siliceous and stand in relief on weathered surfaces. The basal unit consists of olive green, fissile, generally non-calcareous shale. The Flathead Formation is a white to reddish weathering, fine to coarse grained, orthoquartzite that is locally conglomeratic. Beds are thinner and finer grained near the contact with the overlying Silver Hill Formation. Relatively common features include crossbedding, ripple lamination, and locally hematitic cement. Thickness 22-61 m (50-200 ft).
€s	SEDIMENTARY ROCKS OF THE RED LION, HASMARK, SILVER HILL AND FLATHEAD FORMATIONS, UNDIVIDED (CAMBRIAN) The Red Lion Formation is predominately light gray to blue gray limestone with yellow- to red-weathering, wavy, dolomitic and siliceous laminae. Laminae are less common near top of formation. Basal part contains black calcareous shale with minor sandstone and limestone. The thickness is approximately 100 m (330 ft). The Hasmark Formation, approximately 320 m (1,050 ft) thick, is a uniform, light to blue gray dolomite that is separated into an upper and lower part by a recessive shale interval. The lower dolomite is generally darker in color than the upper part and commonly contains oolitic structures and mottled weathered surfaces. The shale interval is as much as 46 m (150 ft) thick and varies from dark brown to reddish purple calcareous shale to reddish limestone and shale. The Silver Hill Formation, approximately 185 m (600 ft) thick, includes calcareous brown, white, and green shale interbedded with laminated limestone underlain by moderately thick-bedded, laminated, light-gray limestone. Laminae are generally siliceous and stand in relief on weathered surfaces. The basal unit consists of olive green, fissile, generally non-calcareous shale. The Flathead Formation is a white to reddish weathering, fine to coarse grained, orthoquartzite that is locally conglomeratic. Beds are thinner and finer grained near the contact with the overlying Silver Hill Formation. Relatively common features include crossbedding, ripple lamination, and locally hematitic cement. Thickness 22-61 m (50-200 ft).
€sm	MARBLE AND QUARTZITE OF THE METAMORPHOSED RED LION, HASMARK, SILVER HILL AND FLATHEAD FORMATIONS (CAMBRIAN) Includes marble, quartzite, gneiss, schist, and phyllite that are the metamorphic equivalents of these formations. This unit is often tectonically thinned. See geologic discussion for more detail.
Ymi	

Ysb	SHEPARD FORMATION (MESOPROTEROZOIC) Dark green siltite and light green argillite in microlaminae and thin beds. Some beds may have a characteristic orange-brown weathering rind. Upper part contains quartzite and siltite. Poorly exposed but weathers into thin platy beds.
Ysn	SNOWSLIP FORMATION (MESOPROTEROZOIC) Mostly red sand to clay couplets with abundant ripples and mud cracks. Siltite-argillite laminae are common near the base. Increasingly argillitic toward top. Some fine- to medium-grained, feldspar-poor quartzite beds are common near the top. Foot thick. The upper portion is mostly flat-laminated, medium-grained, and difficult to distinguish from the Mount Shields Formation. Mud cracks are abundant throughout. In the Anaconda Range are some thin beds of quartzite that contain granule-sized lithic fragments. Thickness is about 37 m (120 ft).
Ysnq	QUARTZITE OF THE METAMORPHOSED SNOWSLIP FORMATION Highly recrystallized quartzite, phyllite, schist, and calc-silicate schist. Metamorphic equivalents of the Snowslip Formation.
Yc	PIEGAN GROUP (MESOPROTEROZOIC) Includes the Wallace and Helena Formations. The upper part, the Helena Formation, is characterized by tan-weathering dolomitic siltite and quartzite. The lower part, the Helena Formation, consists of cycles, from 1 to 9 m (3 to 30 ft) thick, of a basal white quartzite or intraclast unit, overlain by even and wavy bedded argillite without mud cracks, and capped by dolomitic beds. The argillite is difficult to recognize in the typical small outcrop. The unit is made up of parallel, silver-green couplets of darker green siltite and lighter green argillite. It is overlain by beds of tan- or brown-weathering dolomite 30-90 cm (1-3 ft) thick. Mud cracks are common throughout the section. Severe deformation with folding is common. Estimates are problematic, but it is probably at least 1,846 m (6,000 ft) thick.
Ycg	CALC-SILICATE GNEISS OF THE METAMORPHOSED PIEGAN GROUP Greenish, diopside-rich, calc-silicate gneiss, fine-grained quartzite, and schist. Metamorphic equivalent of the Piegan Group.
Yra	RAVALLI GROUP (MESOPROTEROZOIC) Mostly gray, flat-laminated, fine- to medium-grained quartzite and argillite, separated by thin argillite layers. Not significantly metamorphosed. The group is under a gently east-dipping fault; above the fault is Piegan Group. The group is rich in potassium and plagioclase feldspar similar to Mt. Shields. Mud cracks are common nearby exposures of Mount Shields, it contains no pebbles. Soils are common. The unit is tentatively correlated with the Revett Formation. Thickness is at least 914 m (3,000 ft).
Yraq	QUARTZITE OF THE METAMORPHOSED RAVALLI GROUP (MESOPROTEROZOIC) Gray, fine- to medium-grained quartzite in beds 0.15-1.5 m (0.5-5 ft) thick. Phyllite or schist layers. Although original sedimentary structure is obscured, the dominant sediment type appears to be flat-laminated sand, and the argillite. Sediment deformation is common. The unit is tentatively correlated with the Wallace and others (1989, 1992) mapped this unit as Mount Shields. The unit is near Skalkaho Creek and near Storm Lake it can be demonstrated to be equivalent to the overlying Piegan Group. Unit includes a thin zone of locally argillite that separates quartzite from the overlying Piegan Group and may be equivalent to the Formation of the Ravalli Group. Contains 14-21% K-spar and orthoclase. Metamorphism may have affected these concentrations in unmetamorphosed rocks.
Ygg	PELITIC GNEISS AND SCHIST OF THE METAMORPHOSED GREYS (MESOPROTEROZOIC) Mostly reddish-brown-weathering biotite-muscovite schist containing quartz, sillimanite, cordierite, feldspar, garnet, and andalusite. Biotite and cordierite are rare constituents.
Ybg	BIOTITE QUARTZITE OF THE METAMORPHOSED BELT SUPERGROUP Highly recrystallized, fine- to medium-grained quartzite in layers and beds, separated by biotite-rich bands. Although these rocks are almost entirely quartzites, we were unable to identify their equivalent formations.

<p>couplets that are dolomitic and part is red, thinly bedded dolomitic lites. Estimated to be 152 m (500 ft) thick.</p> <p>ud cracks. Green and red, dolomitic amounts of siltite and quartzite up to 1 m thick. Siltite is present in beds less than one meter thick. Siltite to coarse-grained quartzite that is interbedded with mudcracks and mud chips are characteristic of the siltite. Lenticular beds of coarse-grained quartzite are as much as 914 m (3,000 ft) thick.</p> <p>FORMATION (MESOPROTEROZOIC)</p> <p>These rocks that are the metamorphic equivalents of the Wallace Formation, is capped by black argillite in places. They commonly have scoured bases or are overlain by beds of white quartzite clasts in punky, part of the map area. The lower part (0-100 ft) thick and usually incomplete, of lenticular couplets of green siltite and quartzite. However, these cycles are more easily recognized by wavy but not green argillite, by white quartzite, 1-2 m (6-7 ft) thick, and by weathered-out and non-polygonal "crinkle" cracks in this unit makes thickness 100-150 m (300-500 ft) thick.</p> <p>GROUP (MESOPROTEROZOIC)</p> <p>Quartzite, marble, and minor schist.</p> <p>in beds 0.15-1.5 m (0.5-5 ft) thick and is well exposed. Found in the Skalkaho area group. Contains sub-equal amounts of member 2 (unit Yms2), but unlike other units soft sediment deformation is characteristic of the Ravalli Group.</p> <p>PROTEROZOIC)</p> <p>1-5 m (3-15 ft) thick separated by thin siltite. These are only partially preserved, the siltite with crossbeds uncommon. Soft siltite related with the Revett Formation. The siltite fields Formation, but along lower part of the unit to be in stratigraphic contact with siltite. Siltite-bearing pelitic gneiss that is thought to be equivalent to the Saint Regis gneiss. Only 0-10% plagioclase. However, the gneiss is known ways.</p> <p>FORMATION</p> <p>containing variable amounts of kyanite. Kyanite and kyanite pseudomorphs are characteristic.</p> <p>GROUP (MESOPROTEROZOIC)</p> <p>Members 10-30 cm (4-12 in) thick and is well exposed. Most certainly metamorphosed Belt formations.</p>	<p>Range, Montana Bureau of Mines and Geology Bulletin B 84, 54 p., map scale 1:62,500.</p> <p>Elliott, J.E., Waters, M.R., Campbell, W.L., and Avery, D.W., 1984, Mineral resource potential and geologic map of the Dolus Lakes Roadless Area, Powell and Granite Counties, Montana: U.S. Geological Survey Miscellaneous Field Studies Map MF-1640-A, scale 1:50,000.</p> <p>Emmons, W.C., and Calkins, F.C., 1913, Geology and ore deposits of the Philipsburg quadrangle, Montana: U.S. Geological Survey Professional Paper 78, 271 p.</p> <p>Flood, R.E., 1974, Structural geology of the upper Fishtrap Creek area, central Anaconda Range, Deer Lodge County, Montana: Missoula, University of Montana, M.S. thesis, 71 p., map scale 1:62,500.</p> <p>Foster, D.A., Doughty, P.T., Kalakay, T.J., Fanning, C.M., Coyner, S., Grice, W.C., and Vogl, J., 2007, Kinematics and timing of exhumation of metamorphic core complexes along the Lewis and Clark fault zone, northern Rocky Mountains, USA: Geological Society of America Special Paper 434, p. 207-232.</p> <p>Froitzheim, N., Pleuger, J., and Nagel, T.J., 2006, Extraction faults: Journal of Structural Geology, v. 28, p. 1388-1395.</p> <p>Garmezay, Lawrence, and Sutter, J.F., 1983, Mylonitization coincident with uplift in an extensional setting, Bitterroot Range, Montana-Idaho: Geological Society of America Abstracts with Programs, v. 15, p. 578.</p> <p>Grice, W.C., Jr., 2006, Exhumation and cooling history of the middle Eocene Anaconda metamorphic core complex, western Montana: Gainsville, University of Florida, M.S. thesis, 260 p., map scale 1:100,000.</p> <p>Grice, W.C., Jr., Foster, D.A., Kalakay, T.J., Bleick, H.A., and Hodge, K., 2004, Style and timing of crustal attenuation in the Anaconda metamorphic core complex, western Montana: Geological Society of America Abstracts with Programs, v. 36, no. 5, p. 546.</p> <p>Grice, W.C., Jr., Foster, D.A., and Kalakay, T.J., 2005, Quantifying exhumation and cooling of the Eocene Anaconda metamorphic core complex, western Montana: Geological Society of America Abstracts with Programs, v. 37, no. 7, p. 230.</p> <p>Haney, E.M., 2008, Pressure-temperature evolution of metapelites within the Anaconda metamorphic core complex, southwestern Montana: Missoula, University of Montana, M.S. thesis, 110 p.</p> <p>Hawley, K.T., 1974, A study of the mafic rocks along the eastern flank of the Flint Creek range, western Montana: Missoula, University of Montana, M.S. thesis, 39 p., map scale 1:100,000.</p> <p>Hodges, K.V., and Walker, J.D., Extension in the Cretaceous Sevier orogen, North American Cordillera: Geological Society of America Bulletin, v. 104, p. 560-569.</p> <p>Heise, B.A., 1983, Structural geology of the Mount Haggin area, Deer Lodge County, Montana: Missoula, University of Montana, M.S. thesis, 77 p., map scale 1:12,000.</p> <p>Hughes, G.J., Jr., 1970, Precambrian stratigraphy and structure in the Henderson-Willow Creek igneous belt, Granite County, Montana: Houghton, Michigan Tech, M.S. thesis, 93 p., map scale 1:24,000.</p> <p>Hughes, G.J., Jr., 1971, Petrology and tectonic setting of igneous rocks in the Henderson-Willow Creek igneous belt, Granite County, Montana: Houghton, Michigan Tech, Ph.D. dissertation, 236 p.</p> <p>Hyndman, D.W., 1980, Bitterroot dome-Sapphire tectonic block, and example of a plutonic-core gneiss-dome complex with its detached suprastructure: Geological Society of America Memoir 153, p. 427-443.</p>	<p>30' x 60' quadrangle, Montana Bureau of Mines and Geology Open-File Report MBMG 340, 6 p., scale 1:100,000.</p> <p>Lonn, J.D., and C. McDonald, 2004a, Geologic map of the Kelly Lake 7.5' quadrangle, western Montana: Montana Bureau of Mines and Geology Open File Report MBMG 500, 15 p., scale 1:24,000.</p> <p>Lonn, J.D., and McDonald, Catherine, 2004b, Cretaceous(?) syncontractional extension in the Sevier orogen, southwestern Montana: Geological Society of America Abstracts with Programs, v. 36, n. 4, p. 36.</p> <p>Lonn, J.D., Smith, L.N., and McCulloch, R.B., 2007, Geologic map of the Plains 30' x 60' quadrangle, western Montana: Montana Bureau of Mines and Geology Open-File Report MBMG 554, 43 p., scale 1:100,000.</p> <p>Lonn, J.D., McDonald, C., Lewis, R.S., Kalakay, T.J., O'Neill, J.M., Berg, R.B., and Hargrave, P., 2003, Preliminary geologic map of the Philipsburg 30' x 60' quadrangle, western Montana: Montana Bureau of Mines and Geology Open-File Report MBMG 483, 29 p., scale 1:100,000.</p> <p>Lonn, J., and Lewis, R., 2009, Late Cretaceous extension and its relation to the thin stratigraphic section in the Philipsburg area: A field trip to Carpp Ridge: Northwest Geology, v. 38, map scale 1:24,000.</p> <p>Mahorney, J.R., 1956, Geology of the Garrity Hill area, Deer Lodge County, Montana: Bloomington, Indiana University, M.A. thesis, 40 p., map scale 1:25,000.</p> <p>McNulty, Brendan, and Farber, Daniel, 2002, Active detachment faulting above the Peruvian flat slab: Geology, v. 30, p. 567-570.</p> <p>Marvin, R.F., Mehnert, H.H., Naeser, C.W., and Zortman, R.E., 1989, USGS radiometric ages--compilation "c", part 5: Colorado, Montana, Utah, and Wyoming: IsochronWest, v. 53, p. 14-19.</p> <p>McGill, G.E., 1961, Geologic map of the northwest flank of the Flint Creek Range, western Montana: Montana Bureau of Mines and Geology Geologic Map GM 3, scale 1:32,000.</p> <p>Mutch, T.A., 1961, Geologic map of the northeast flank of the Flint Creek Range, western Montana: Montana Bureau of Mines and Geology Special Publication SP 22, scale 1:62,500.</p> <p>Noel, J.A., 1956, The geology of the east end of the Anaconda Range and adjacent areas, Montana: Bloomington, Indiana University, Ph.D. dissertation, 74 p.</p> <p>O'Connell, M.P., 2001, Geometry, kinematics, and emplacement mechanisms of the Philipsburg Batholith within the Sevier fold and thrust belt, Flint Creek Range, western Montana: Bozeman, Montana State University, M.S. thesis, 91 p., map scale 1:24,000.</p> <p>O'Neill, J.M., 2005, Syntectonic Anaconda Conglomerate (new name)—A stratigraphic record of early Tertiary brittle-ductile extension and uplift in southwestern Montana: U.S. Geological Survey Professional Paper 1700, Chapter A, p. 1-15.</p> <p>O'Neill, J.M., Lonn, J.D., and Kalakay, T., 2002, Early Tertiary Anaconda Metamorphic Core Complex, southwestern Montana: Geological Society of America Abstracts with Programs, v. 34, no. 4, p. A10.</p> <p>O'Neill, J.M., Lonn, J.D., Lageson, D.R., and Kunk, M.J., 2004, Early Tertiary Anaconda metamorphic core complex, southwestern Montana: Canadian Journal of Earth Sciences, v. 41, p. 63-72.</p> <p>Pederson, R.J., 1976, Geology of the upper Rock Creek drainage, Granite County, Montana: Butte, Montana College of Mineral Science and Technology, M.S. thesis, 238 p., map scale 1:24,000.</p>	<p>Winston, Don, 1986c, Sedimentation and tectonic of the middle Proterozoic Belt Basin, and their influence on Phanerozoic compression and extension in western Montana and northern Idaho, in Peterson, J.A., ed., Paleotectonics and Sedimentation in the Rocky Mountain Region, United States: American Association of Petroleum Geologists Memoir 41, p. 87-118.</p> <p>Winston, Don, 1998, Lacustrine cycles in the Helena and Wallace Formations, middle Proterozoic Belt Supergroup, western Montana, in Berg, R.B., ed., Proceedings of Belt Symposium III: Montana Bureau of Mines and Geology Special Publication SP 112, p. 70-86.</p> <p>Winston, Don, McGee, D., and Quattlebaum, D., 1986, Stratigraphy and sedimentology of the Bonner Formation, middle Proterozoic Belt Supergroup, western Montana, in Roberts, S.M., Belt Supergroup: A guide to Proterozoic rocks of western Montana and adjacent areas: Montana Bureau of Mines and Geology Special Publication SP 94, p. 183-195.</p> <p>Winston, Don, and Wallace, C.A., 1983, The Helena Formation and the Missoula Group at Flint Creek Hill, near Georgetown Lake, western Montana, in Hobbs, S.W., ed., Guide to Field Trips, Belt Symposium II: Department of Geology, University of Montana, Missoula, p. 66-81.</p>
--	--	--	--

TKa	GRANITIC ROCKS (EARLY TERTIARY AND LATE CRETACEOUS)
TKgd	Non-foliated biotite-muscovite monzogranite, leucomonzogranite, and granodiorite.
TKgd	GRANODIORITIC ROCKS (EARLY TERTIARY AND LATE CRETACEOUS)
TKgb	Non-foliated biotite granodiorite, hornblende-biotite granodiorite, tonalite, and quartz diorite.
TKgb	GABBROIC ROCKS (EARLY TERTIARY AND LATE CRETACEOUS)
TKps	Gabbro, microgabbro, diorite, and lamprophyre.
TKps	PYROXENITE AND SYENITE (EARLY TERTIARY OR LATE CRETACEOUS)
Kqd	Pyroxenite and syenite found near the western boundary of the map area.

Ymiq	MISSOULA GROUP, UNDIVIDED (MIDDLE PROTEROZOIC)
	Includes, in descending order, the Garnet Range, McNamara, Bonner, Mount Shields, Shepard, and Snowslip Formations. Total thickness as much as 3,261 m (10,700 ft).
Ymiq	QUARTZITE AND PHYLLITE OF THE METAMORPHOSED MISSOULA GROUP (MIDDLE PROTEROZOIC)
Ygr	Quartzite, phyllite, schist, calc-silicate rocks, and gneiss that are metamorphic equivalents of the Missoula Group. The Bonner, McNamara, and Garnet Range Formations have been eroded off. In addition, this unit is often tectonically thinned, and varies from 62 to 677 m (200 to 2,200 ft) in thickness. See the geologic discussion for more detail.

Ybg	GNEISS, QUARTZITE, CALC-SILICATE ROCKS, SCHIST, AND PHYLLITE OF THE BELT SUPERGROUP (MESOPROTEROZOIC)
	Highly recrystallized gneissic and schistose rocks to which we assign group equivalents, but that probably belong to the Belt Super

Montana Bureau of Mines and Geology
Open File 483, Plate 2

Geologic Map of the Philipsburg 30' x 60' Quadrangle, Western Montana

Jeffrey D. Lonn¹, Catherine McDonald¹, Reed S. Lewis²,
Thomas J. Kalakay³, J. Michael O'Neill⁴,
Richard B. Berg¹, and Phyllis Hargrave¹

2003 (Revised 2010)

¹ Montana Bureau of Mines and Geology; ² Idaho Geological Survey;
³ Rocky Mountain College; ⁴ U.S. Geological Survey

GEOLOGIC SUMMARY

Introduction

Montana Bureau of Mines and Geology's new Geologic Map of the Philipsburg 30' x 60' Quadrangle represents a revised version of the Preliminary Geologic Map of the Philipsburg Quadrangle (Lonn and others, 2003) based on new field work by Lonn and Lewis from 2003 to 2008. This new field work addressed structural and stratigraphic problems revealed by, but not resolved on, the previous map.

Structural Geology

The Philipsburg quadrangle can be divided into two major structural domains separated by the north-northeast-striking Georgetown-Philipsburg thrust system (fig. 1). The eastern structural domain, comprising the Flint Creek and northeastern Anaconda Ranges, is characterized by upper greenschist to upper amphibolite facies metamorphism, tight folds, closely spaced faults, and a complex structural history. The western domain, previously termed the Sapphire tectonic block (Hyndman and others, 1975) or Skalkaho slab (Doughty and Sheriff, 1992), is an allochthon composed mostly of low-grade metasedimentary rocks deformed into upright, open folds and cut by numerous reverse and normal faults. Both domains are extensively intruded by late Cretaceous to early Tertiary granitic and dioritic plutons.

Eastern Structural Domain

Structural geology is extremely complex within the eastern domain. Major east-directed thrust faults, represented by the Georgetown-Philipsburg thrust system, presumably buried the rocks of the eastern domain to mid-crustal depths in late Cretaceous time. An increase in metamorphic grade from west to east probably reflects greater uplift in the east. The southeastern-most part of the Anaconda Range contains relict kyanite and kyanite pseudomorphs (Kalakay and others, 2003; Grice, 2006) indicative of high-pressure metamorphism, overprinted by a high-temperature, lower pressure metamorphic event at about 80-75 Ma (Grice and others, 2004, 2005; Grice, 2006; Haney, 2008).

In the Flint Creek and Anaconda Ranges, the Mesoproterozoic through Mesozoic metasedimentary sequence appears to be tectonically attenuated by an array of bedding-parallel fabrics and structures that include concordant mylonitic shear zones that cut out stratigraphic section, zones of vertical shortening that flatten the units through pure shear and plastic flow, and brittle bedding-parallel faults that place younger units over older units (Lonn and McDonald, 2004a,b; Lonn and Lewis, 2009). Parallel solid-state fabrics are present in the oldest (> 75 Ma) late Cretaceous plutons (units Kgdf, Kqdf) intruding the metasediments (Hawley, 1974; Desmarais, 1983; Grice and others, 2005;

Grice, 2006). The strain fabrics apparently formed during the 75-80 Ma high temperature metamorphic event (Grice and others, 2004, 2005; Grice, 2006), and they have been deformed with the beds into tight, NNE-trending, west-verging, asymmetric to overturned folds whose east-dipping axial planes appear to become gently inclined with increasing structural depth. Undeformed late Cretaceous to early Tertiary plutons intrude the metasediments. Most plutons are sheet-like and roughly concordant to bedding, and their intrusion may have been synchronous with the folding.

The eastern flanks of the Anaconda and Flint Creek Ranges are overprinted by structures and fabrics associated with the Eocene Anaconda metamorphic core complex (O'Neill and others, 2002, 2004). The confusing geology in the southeastern corner of the Philipsburg quadrangle typifies map patterns in this extensional terrane. Here, chaotic, brecciated sedimentary rocks of the upper plate are separated from lower plate metamorphic and plutonic rocks by brittle detachment faults and a greenschist-facies mylonitic shear zone. The sinuous and discontinuous Anaconda mylonite, gently folded and broken by faults, extends for more than 100 km (62 miles) along strike. It dips gently east, has a top-to-the-east shear sense, and was active from at least 53 to 47 Ma (Grice and others, 2005; Grice, 2006). Mineral lineations consistently plunge gently ESE (102o-108o), bearings almost identical to those associated with the coeval (Foster and others, 2007) Bitterroot metamorphic core complex 100 km (62 miles) to the west and outside the quadrangle.

The Georgetown-Philipsburg Thrust

The Georgetown-Philipsburg thrust system divides the western and eastern domains. It is a complex imbricate fault system that places Mesoproterozoic Piegan Group of the Belt Supergroup over upper Paleozoic and Mesozoic sediments for a total stratigraphic separation of 7,400 m (24,000 ft). Regional cross sections that restore the slight angular unconformity at the Belt-Cambrian contact suggest about 35 km (22 miles) of horizontal displacement. The Georgetown fault is folded, perhaps by the same folds that deform the rocks of the eastern domain, and the thrust is also overprinted by normal faults that obscure the original thrust geometries along much of its trace. A minimum age of 78 Ma for the fault is inferred from cross-cutting late Cretaceous plutons (Hyndman and others, 1982; Desmarais, 1983; Marvin and others, 1989; Wallace and others, 1992).

Western Structural Domain

West of the Georgetown-Philipsburg thrust is the Sapphire allochthon, mostly composed of gently folded, low-grade, Mesoproterozoic Belt Supergroup rocks intruded by late Cretaceous to early Tertiary plutons. However, the Sapphire allochthon is clearly not an intact block. It is complexly deformed by faults and shear zones of several types: 1) major

were unable to assign formation or group.

Hyndman, D.W., 1983, The Idaho Batholith and associated plutons, Idaho and western Montana, in Roddick, J.A., ed., Circum-Pacific plutonic terranes: Geological Society of America Memoir 159, p. 213-240.

Hyndman, D.W., Obradovich, J.D., and Ehinger, R., 1972, Potassium-argon age determinations of the Philipsburg Batholith: Geological Society of America Bulletin, v. 83, p. 473-474.

Hyndman, D.W., Talbot, J.L., and Chase, R.B., 1975, Boulder Batholith: a result of emplacement of a block detached from the Idaho Batholith infrastructure?: Geology, v.3, p. 401-404.

Poulter, G.J., 1956, Geology of the Georgetown thrust area southwest of Philipsburg, Montana: Princeton, NJ, Princeton University Ph.D. dissertation, map scale 1:24,000.

Presley, M.W., 1971, Igneous and metamorphic geology of the Willow Creek drainage basin, southern Sapphire Mountains, Montana: Missoula, University of Montana, M.S. thesis, 64 p., map scale 1:31,200.

Prinz, W.C., 1967, Geology and ore deposits of the Philipsburg district, Granite County, Montana: U.S. Geological Survey Bulletin 1237, 66 p., map scale 1:16,000.

reverse faults like the Stony Lake thrust; 2) complex anastomosing fault systems that contain both reverse and normal faults, represented by the Ranch Creek fault system; 3) the NE-striking, near vertical, locally mylonitic Skalkaho shear zone; 4) low-angle younger-over-older normal(?) faults, exemplified by the bedding-parallel Shadow Lake and Burnt Fork detachments, and by the discordant Railroad Creek detachment; 5) high-angle normal(?) faults such as the Daly Creek fault and basin-bounding upper Willow Creek fault. In addition, an enigmatic area of kyanite-bearing, Belt-protolith metamorphic rock exists near the western border of the map in the upper plate of the Bitterroot metamorphic core complex. The gently east-dipping Bitterroot detachment fault defines the western boundary of the Sapphire allochthon and is located just west of the map area. The southwestern corner of the map is composed mostly of late Cretaceous to early Tertiary granitic plutons that post-date all but the high-angle normal(?) faults. On the east slope of the Sapphire Range is the rhyolitic Eocene(?) Rock Creek volcanic field, the probable source of the sapphires (Berg and Dahy, 2002) for which the mountains are named.

Stratigraphy

Middle Proterozoic Belt Supergroup sedimentary and metasedimentary rocks dominate the map area, and Belt nomenclature from type localities in northwestern Montana has been applied to these rocks. Of particular significance is the affirmation of Ravalli Group rocks (Revett Formation?) in the Anaconda Range and southern Sapphire Range. These rocks bring the exposed Belt section thickness to more than 6,100 m (20,000 ft) in the western part of the Philipsburg quadrangle, not including possible lower Belt rocks buried beneath the surface. However, in the eastern domain, as discussed above, the thickness of the Belt section has been thinned by a combination of pre-middle Cambrian erosion and Cretaceous tectonism to less than 4,000 m (13,000 ft), even though the entire lower Belt through Missoula Group section is present. The Paleozoic and Mesozoic sections also appear to have been tectonically attenuated in the eastern domain; they have been eroded off the western domain.

Missoula Group rocks appear to coarsen toward the southern boundary of the Belt Basin that lies only 50 km (30 miles) south of the Philipsburg quadrangle. Both the Mount Shields member 2 and the Bonner Formation contain significant conglomerate in the Anaconda and southern Sapphire Ranges. The McNamara Formation, an argillite-rich unit at its type locality near Missoula, becomes a medium-grained feldspathic arenite in the southern Sapphire Range and near Georgetown Lake. In contrast, Ravalli Group quartzites do not display similar southward-coarsening trends.

Metamorphic Rocks

Amphibolite facies regional metamorphic rocks are common in the quadrangle, although previous maps identified them only as their sedimentary equivalents. Because the distribution of metamorphic rocks is so important to interpreting the structural geology, we have attempted to show them on the map as metamorphic equivalents of the various units. Metamorphism probably occurred prior to intrusion of most of the major plutons in the Philipsburg quadrangle (Stuart, 1966; Grice, 2006; Haney, 2008).

In addition, areas of mylonitic foliation are shown along the eastern flank of the Anaconda metamorphic core complex. Areas of significant tectonic breccia are also shown.

Regional Structural Interpretation

The earliest tectonic event that can be documented in the region is gentle (2° -5°) westward tilting and subsequent erosion of the Mesoproterozoic Belt Supergroup before deposition of the middle Cambrian Flathead Formation. Although there are some disconformities present within the Paleozoic and Mesozoic stratigraphic sections, no other major tectonic events can be identified until the start of the Cretaceous Sevier orogeny. During Sevier orogenesis, east-directed thrust systems like the Stony Lake, Ranch Creek, and Georgetown-Philipsburg thickened the crust and buried the footwall rocks (the eastern domain) to mid-crustal depths beneath the rocks of the western domain. The footwall rocks then underwent high-pressure metamorphism followed by high-temperature, low-pressure metamorphism that coincided with the bedding-parallel fabrics that are associated with the tectonically attenuated stratigraphic section (Kalakay and others, 2003; Grice, 2006). The thinning of the entire >12,200-meter-thick (40,000 ft) metasedimentary section, the faults and shear zones that always omit and never duplicate section, and the dominance of pure shear (coaxial strain) fabrics over simple shear (non-coaxial strain) fabrics suggest to us that the thin stratigraphic section and bedding-parallel fabrics resulted from a period of synorogenic, late Cretaceous extension that occurred in a convergent tectonic setting synchronously with thrusting in the foreland to the east. In fact, there is evidence that some thrusting in the Philipsburg region was coeval with or postdates this extension: 1) detachment faults are duplicated by later thrusts (Lonn and McDonald, 2004a); and 2) similar bedding-parallel faults omit the Snowslip and Shepard Formations on parts of both the hanging wall and footwall of the Georgetown thrust. Hodges and Walker (1992) cite extensive evidence for similar late Cretaceous extension synchronous with thrusting in other areas of the Sevier hinterland, while numerous studies in the Andes and Himalaya have documented the occurrence of active extension in a convergent setting (Dalmayrac and Molnar, 1981; Burchfiel and

Royden, 1985; McNulty and Farber, 2002). The postulated late Cretaceous extension may have been facilitated by thermal heating that resulted from crustal thickening and the emplacement of the earliest plutons; in turn, the extension may have generated more plutonism, represented by the voluminous 75-60 Ma intrusions, through decompression melting.

Although some folding undoubtedly occurred during thrusting, the puzzling, west-vergent folds formed during or after most of the thrusting and the proposed extensional structures. The folds may be synchronous with many of the sheet-like, 75-65 Ma intrusions. These folds that verge west--the wrong way--have been difficult to explain. Although they have been attributed to thin-skinned thrust tectonics (Emmons and Calkins, 1913; Csejtey, 1962; Flood, 1974; Wallace and others, 1992), they may represent hot, ductile middle crust (infrastructure) that continued to plastically deform beneath the brittle, cold, upper crust (superstructure) after deformation in the superstructure had ceased (Culshaw and others, 2006).

Convergent tectonism in the region ended in the Paleocene (Harlan and others, 1988) and was immediately followed by crustal extension represented by the Eocene Anaconda metamorphic core complex (O'Neill and others, 2002, 2004). The main Anaconda detachment initiated at about 53 Ma and the mylonitic shear zone was active until at least 47 Ma, and possibly until 30 Ma (Grice and others, 2004; Grice, 2006). The Bitterroot metamorphic core complex just beyond the western border of the Philipsburg quadrangle developed at the same time, and the two are thought to be "nested" core complexes (Foster and others, 2007). Eocene Lowland Creek volcanic rocks (unit Tlc) interfinger with coarse clastic and landslide deposits of the Anaconda beds (unit Tac) that were derived from unroofing of the Anaconda core complex (O'Neill and others, 2004; O'Neill, 2005). Rhyolitic rocks of the Rock Creek volcanic field are probably also of Eocene age.

Most high-angle and listric normal faults appear to be Eocene and younger. Some bound Tertiary valleys like the Upper Willow Creek Valley. Others, like those of the Ranch Creek fault zone and the Georgetown thrust zone, merge and anastomose with reverse faults, and are thought to represent normal-sense reactivation of thrust faults (Lewis, 1998b), although some could have formed synchronously with the thrusts through a constructional strain/extrusion process (Reid and others, 1995; Froitzheim and others, 2006; Lonn and others, 2007). Voluminous sedimentary deposits (units Ts, Tac, Taf) filled basins developed by the Tertiary normal faults.