

CORRELATION OF MAP UNITS

Quaternary	Quaternary	Quaternary	Quaternary
Pleistocene	Pleistocene	Pleistocene	Pleistocene
Pliocene	Pliocene	Pliocene	Pliocene
Upper Cretaceous	Upper Cretaceous	Upper Cretaceous	Upper Cretaceous
Lower Cretaceous	Lower Cretaceous	Lower Cretaceous	Lower Cretaceous
Upper Jurassic	Upper Jurassic	Upper Jurassic	Upper Jurassic
Middle Jurassic	Middle Jurassic	Middle Jurassic	Middle Jurassic
Lower Jurassic	Lower Jurassic	Lower Jurassic	Lower Jurassic
Permian	Permian	Permian	Permian
Pennsylvanian	Pennsylvanian	Pennsylvanian	Pennsylvanian
Upper Mississippian	Upper Mississippian	Upper Mississippian	Upper Mississippian
Middle Mississippian	Middle Mississippian	Middle Mississippian	Middle Mississippian
Lower Mississippian	Lower Mississippian	Lower Mississippian	Lower Mississippian
Devonian	Devonian	Devonian	Devonian
Middle Devonian	Middle Devonian	Middle Devonian	Middle Devonian
Middle and Upper Cambrian			
Archean	Archean	Archean	Archean

DESCRIPTION OF MAP UNITS

Quaternary	Aluvium (Holocene) : Gravel, sand, silt, and clay along active channels and floodplains. Deposits are locally extensive in tributaries draining areas underlain by Cretaceous sandstone, and shale bedrock is sand, silt, and clay.	Bearpaw Shale (Upper Cretaceous) : Dark-gray shale, brownish-gray calcareous concretions and nodules are common. Part of formation contains thin, well-sorted, thin-bedded, micaceous, fine to coarse-grained sandstone. The thickness is 200 to 800 feet, thinning to the west.	Kootenai Formation (Lower Cretaceous) : Mostly reddish-brown to gray, micaceous, fine to coarse-grained sandstone. The thickness is 200 to 250 feet.
Pleistocene	Colluvium (Holocene and Pleistocene) : Locally derived thin veneer concealing bedrock, but locally as thick as 20 feet. Commonly grades into Qal. Locals are well-sorted, rounded cobbles derived from alluvial terrace gravel.	Judith River Formation (Upper Cretaceous) : Interbedded brownish-gray calcareous sandstone and shale. Locally contains a massive, off-forming sandstone, representing thin-bedded, micaceous, fine to coarse-grained sandstone. The thickness is 200 to 800 feet, thinning to the west.	Morrison Formation (Upper Jurassic) : Variegated, mainly reddish-brown to gray, micaceous, fine to coarse-grained sandstone. The thickness is 200 to 250 feet.
Pliocene	Alluvial fan deposits (Holocene and Pleistocene) : Gravel, sand, silt, and clay deposited in fans formed by modern streams along major valley margins. Display characteristic fan-shaped topography. Typically grade upstream into Qal. Thicknesses range from very thin to as much as 50 feet at heads of fans.	Eagle Sandstone (Upper Cretaceous) : Light brownish-gray to very pale, orange very fine to fine-grained, cross-bedded sandstone. The thickness is 200 to 800 feet, thinning to the west.	Ellis Group, undivided
Upper Cretaceous	Older alluvial fan deposits (Holocene?) : Deposits similar to Qal but overlain by Qal and are erosionally dissected; occur only along the southwest margin of the Pryor Mountains.	Telegraph Creek Formation (Upper Cretaceous) : Dark-gray to black, micaceous, fine to coarse-grained sandstone. The thickness is 200 to 800 feet, thinning to the west.	Swain Formation (Upper Jurassic) : Interbedded medium-gray shale, limestone and limy sandstone. The thickness is 200 to 250 feet.
Lower Cretaceous	Landslide deposits (Holocene and Pleistocene) : Unconsolidated mixture of soil and blocks of bedrock. Deposits are locally extensive in tributaries draining areas underlain by Cretaceous sandstone, and shale bedrock is sand, silt, and clay.	Niobrara Shale (Upper Cretaceous) : Olive-gray to dark brownish-gray, micaceous, fine to coarse-grained sandstone. The thickness is 200 to 800 feet, thinning to the west.	Riverton Formation (Upper Jurassic) : Mostly pale greenish-gray very fossiliferous shale with minor interbedded reddish-brown limestone. The thickness is 200 to 250 feet.
Upper Jurassic	Pediment surface deposits (Pleistocene) : Deposits of sand, silt, and clay mixed with cobble-size clasts of locally derived sandstone. Deposits are locally extensive in tributaries draining areas underlain by Cretaceous sandstone, and shale bedrock is sand, silt, and clay.	Frontier Formation (Upper Cretaceous) : Light brownish-gray to very pale, micaceous, fine to coarse-grained sandstone. The thickness is 200 to 800 feet, thinning to the west.	Piper Formation (Middle Jurassic) : Interbedded medium gray and pale reddish-gray, thin-bedded limestone and medium-gray shale. Includes thin, interbedded gypsum. The thickness is 200 to 250 feet.
Middle Jurassic	Landslide deposits (Holocene and Pleistocene?) : Unconsolidated mixture of soil and blocks of bedrock along the flanks of the Pryor Mountains south. Much of surface form is obliterated by later erosion.	Greenhorn Formation (Upper Cretaceous) : Shale, dark brownish-gray to black, micaceous, fine to coarse-grained sandstone. The thickness is 200 to 800 feet, thinning to the west.	Chugwater Formation (Lower Triassic) : Interbedded medium gray and pale reddish-gray, thin-bedded limestone and medium-gray shale. Includes thin, interbedded gypsum. The thickness is 200 to 250 feet.
Lower Jurassic	Pediment gravel deposits (Holocene and Pleistocene?) : Gravel and sand deposited on pediment surfaces. Deposits are locally extensive in tributaries draining areas underlain by Cretaceous sandstone, and shale bedrock is sand, silt, and clay.	Mowry Shale (Upper Cretaceous) : Interbedded siliceous, very fine to fine-grained sandstone, siltstone, and shale. The thickness is 200 to 800 feet, thinning to the west.	Anderson Formation (Lower Pennsylvanian and Upper Mississippian) : Interbedded grayish-gray to light-red mudstone, limestone, and siltstone. Limestones are commonly cherty. The thickness is 200 to 250 feet.
Permian	Calcareous tufa (Pleistocene and Pliocene?) : Light-gray and grayish-brown, vuggy, limestone tufa deposits. Deposits are locally extensive in tributaries draining areas underlain by Cretaceous sandstone, and shale bedrock is sand, silt, and clay.	Thermopsis Shale (Upper Cretaceous) : Interbedded siliceous, very fine to fine-grained sandstone, siltstone, and shale. The thickness is 200 to 800 feet, thinning to the west.	Madison Group, undivided (Middle Mississippian) : Limestone and siliceous limestone. The thickness is 200 to 250 feet.
Pennsylvanian	Alluvial terrace gravel (Pleistocene and Pliocene?) : Light-gray and grayish-brown, vuggy, limestone tufa deposits. Deposits are locally extensive in tributaries draining areas underlain by Cretaceous sandstone, and shale bedrock is sand, silt, and clay.	Lance Formation (Upper Cretaceous) : Interbedded light brownish-gray, silt and ledge-forming, fine-grained, thick-bedded, micaceous, fine to coarse-grained sandstone. The thickness is 200 to 800 feet, thinning to the west.	Jefferson Formation (Upper Devonian) : Limestone, light brownish-gray, locally exposed. The thickness is 200 to 250 feet.
Upper Mississippian	Reversal fault : Open teeth on upthrown block	Labo Member, Fort Union Formation (Pliocene) : Primarily dark gray to olive shale, locally yellowish-gray. The thickness is 200 to 250 feet.	Bighorn Dolomite (Middle Devonian) : Dolomite and dolomitic limestone, very light gray to very pale-orange. The thickness is 200 to 250 feet.
Middle Mississippian	Strike and dip of inclined beds	Tullock Member, Fort Union Formation (Pliocene) : Yellowish-gray, fine to medium-grained, ledge-forming sandstone, cross-bedded in part. The thickness is 200 to 250 feet.	Cambrian Sedimentary rocks undivided (Middle and Upper Cambrian) : Limestone, light brownish-gray, locally exposed. The thickness is 200 to 250 feet.
Lower Mississippian	Strike and dip of overturned beds	Labo Member, Fort Union Formation (Pliocene) : Primarily dark gray to olive shale, locally yellowish-gray. The thickness is 200 to 250 feet.	Granitic gneiss, schist, and biotite schist (Archean) : Pale to moderate red gneiss, granite, medium-dark-gray quartz-feldspathic gneiss, biotite schist, and biotite schist. The thickness is 200 to 250 feet.
Devonian	Anticline : Showing trace of axial plane and direction of plunge, dashed where approximately located, dotted where concealed	Labo Member, Fort Union Formation (Pliocene) : Primarily dark gray to olive shale, locally yellowish-gray. The thickness is 200 to 250 feet.	
Middle Devonian	Syncline : Showing trace of axial plane and direction of plunge, dotted where concealed	Labo Member, Fort Union Formation (Pliocene) : Primarily dark gray to olive shale, locally yellowish-gray. The thickness is 200 to 250 feet.	
Middle and Upper Cambrian	Overturned syncline : Showing trace of axial plane and direction of dip of bedding, dashed where approximately located, dotted where concealed	Labo Member, Fort Union Formation (Pliocene) : Primarily dark gray to olive shale, locally yellowish-gray. The thickness is 200 to 250 feet.	
Archean	Monocline : Showing axial plane trace of anticlinal structure, dashed where approximately located, dotted where concealed; shorter arrow on more steeply dipping limb	Labo Member, Fort Union Formation (Pliocene) : Primarily dark gray to olive shale, locally yellowish-gray. The thickness is 200 to 250 feet.	

MAP SYMBOLS

- Dashed line: Contact dashed where approximately located, dotted where concealed
- Dotted line: Fault: Dashed where approximately located, dotted where concealed
- Open teeth: Reverse fault: Open teeth on upthrown block
- Horizontal line: Strike and dip of inclined beds
- Arrow: Strike and dip of overturned beds
- Red dashed line: Anticline: Showing trace of axial plane and direction of plunge, dashed where approximately located, dotted where concealed
- Red dotted line: Syncline: Showing trace of axial plane and direction of plunge, dotted where concealed
- Red dashed line with arrow: Overturned syncline: Showing trace of axial plane and direction of dip of bedding, dashed where approximately located, dotted where concealed
- Red dashed line with arrow: Monocline: Showing axial plane trace of anticlinal structure, dashed where approximately located, dotted where concealed; shorter arrow on more steeply dipping limb
- Blue line: Water

INDEX OF PREVIOUS GEOLOGIC MAPPING

10	1974h
11	1974i
12	1974j
13	1974k
14	1974l
15	1974m
16	1974n
17	1974o
18	1974p
19	1974q

REFERENCES

Andrews, D.A., Prens, W.G., and Eagle, D.H.: 1947. Geologic map of the Big Horn Basin, Wyoming and Montana, showing terrace deposits and physiographic features. U.S. Geological Survey, U.S. Geol. Investigation Bulletin 66, scale 1:250,000.

Blackstone, D.L.: 1974a. Preliminary geologic map of the Deep Creek SE quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 67, scale 1:24,000.

Blackstone, D.L.: 1974b. Preliminary geologic map of the Deep Creek SE quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 67, scale 1:24,000.

Blackstone, D.L.: 1974c. Preliminary geologic map of the Castle Rock quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 68, scale 1:24,000.

Blackstone, D.L.: 1974d. Preliminary geologic map of the Big Ice Cave quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 69, scale 1:24,000.

Blackstone, D.L.: 1974e. Preliminary geologic map of the Bear Canyon quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 66, scale 1:24,000.

Blackstone, D.L.: 1974f. Preliminary geologic map of the Bowler Creek SE quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 67, scale 1:24,000.

Blackstone, D.L.: 1974g. Preliminary geologic map of the Red Pryor Mountain quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 68, scale 1:24,000.

Blackstone, D.L.: 1974h. Preliminary geologic map of the Big Ice Cave quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 69, scale 1:24,000.

Blackstone, D.L.: 1974i. Preliminary geologic map of the Section House Draw quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 69, scale 1:24,000.

Blackstone, D.L.: 1974j. Preliminary geologic map of the Mystery Creek quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 70, scale 1:24,000.

Blackstone, D.L.: 1974k. Preliminary geologic map of the Indian Spring quadrangle, Big Horn County, Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 71, scale 1:24,000.

Blackstone, D.L.: 1974l. Preliminary geologic map of the East Pryor Mountain quadrangle, Carbon County, Montana. Montana Bureau of Mines and Geology Special Publication 69, scale 1:24,000.

Campbell, N.P.: 1978. Caves of Montana. Montana Bureau of Mines and Geology Bulletin 105, 169 p.

Boett, G.A., and Wilcox, R.E.: 1950. Map showing localities and mineral distribution of the Hudson River, Meigs Falls, and Judith Creek ash beds (Hemlock Ash Beds) of Pliocene and Pleistocene age in western United States and southern Canada. U.S. Geological Survey Miscellaneous Investigations Series Map I-1325, scale 1:4,000,000.

Knapp, R.S., and Moulton, G.F.: 1931. Geology and mineral resources of parts of Carbon, Big Horn, Yellowstone, and Stillwater counties. U.S. Geological Survey Bulletin 824-A, 70 p., pl. 1, scale 1:250,000.

Lopez, D.A.: 1995. Field guide to the northern Pryor-Bighorn structural block, south central Montana. Montana Bureau of Mines and Geology Open-File Report MBMG 82, scale 1:24,000.

Lopez, D.A.: 1996. Structural geology of the northern Pryor-Bighorn uplift, south central Montana [abs.]. American Association of Petroleum Geologists Bulletin, v. 80, p. 914-915.

Maskin, J.H.: 1937. Erosional history of the Big Horn Basin. Wyoming Geological Society of America Bulletin, v. 48, p. 813-894.

McCormell, D.A.: 1994. Five-finger, basement-involved fault propagation basin and gas investigations. Wyoming Geological Society of America Bulletin, v. 48, p. 813-894.

Rehms, J.C.: 1985. Evidence for Quaternary tectonism in the northern Big Horn Basin, Wyoming and Montana. Geology, v. 13, p. 364-367.

Richards, P.W.: 1955. Geology of the Bighorn Canyon-Hurdin area, Montana and Wyoming. U.S. Geological Survey Bulletin 1030, 83 p., scale 1:62,500.


Richards, P.W., and Rogers, C.F.: 1951. Geology of the Hardin area, Big Horn and Yellowstone counties, Montana. U.S. Geological Survey Bulletin 804-QM-111, sheet 1, scale 1:50,000.

Sarberg, C.A.: 1961. Distribution and thickness of Devonian rocks in the Big Horn Basin and northern Wyoming. U.S. Geological Survey Bulletin 1115-D, 127 p.

Stewart, J.C.: 1974. Geology of the Dryhead-Garvin basin, Big Horn and Carbon counties, Montana. Montana Bureau of Mines and Geology Special Publication 17, scale 1:62,500.

Thom, W.T., Jr., Hall, G.M., Wegmann, C.H., and Moulton, G.F.: 1935. Geology of Big Horn County and the Crow Indian Reservation, Montana. U.S. Geological Survey Bulletin 806, 200 p.

Zimmerman, E.A.: 1960. Geology of the Bluewater Springs area, Carbon County, Montana. U.S. Geological Survey Water-Supply Paper 1779-A, 24 p.



**Geologic Map of the
Bridger 30' x 60' Quadrangle, Montana**
by
David A. Lopez
2000



Map not to be construed from
Publication Office
1100 West Park Street, Butte, Montana 59717-0997
Phone: (406) 338-4200
Fax: (406) 338-4411
http://www.mbg.mt.edu

UTM grid coordinates (GCS) and UTM projection
at center of map
Digital reproduction